THE EPHRAIM LIBRARY

3 1151 02676 8162

Library

Johns Hopkins University

Investigations on the Glare Quatic.

By the author

Submitted to the Board of University Studies of the
Johns Hopkins University in accordance with the require-
ments for the degree of Doctor of Philosophy.

	Page
1. Introduction	1
2. The Undulation	4
3. The Discriminant of the Hessian	5
4. The Discriminant of S, T, H	10
5. The Discriminant of t, f	13
6. The Twenty-one Lines	15
7. The Invariant of unknown Degree	17
8. Certain Other Invariants	19
9. The Eliminant of S, T, H .	25
10. The Eliminant of the Polar Cubic, Conic, and Line.	29
11. The Polar Conic of Two Points	32
12. Salmon's Cones	36
Notes	49

§ 1. Introduction.

The plane quartic $(xy)^4$ is taken in the form

$$\begin{aligned} & ax_0^4 \\ & + 4a_1x_0^3x_1 + 4a_2x_0^3x_2 \\ & + 6hx_0^2x_1^2 + 12lx_0^2x_1x_2 + 6gx_0^2x_2^2 \\ & + 4b_0x_0x_1^3 + 12mx_0x_1^2x_2 + 12nx_0x_1x_2^2 + 4c_0x_0x_2^3 \\ & + 4b_1x_1^4 + 4b_2x_1^3x_2 + 6fx_1^2x_2^2 + 4c_1x_1x_2^3 + 4x_2^4. \end{aligned}$$

It will be convenient first to mention briefly certain well-known forms connected with it that will be made use of in this article.

Of these several arise from the polar forms $(xy)^3(xy)$, $(xy)^2(xy)^2$, $(xy)(xy)^3$. Since of each of these there is an D^2 , the placing of one, two, or three conditions on the curves they represent results, respectively, in a locus for the pole, or a set of points, and in an invariant condition to be satisfied by the quartic.

Upon the polar line the only condition placeable is that it vanish identically. This, as is well known, means

that the quartic has a double point, it requires the vanishing of its discriminant, an A^{27} .^{*}

The polar cubic may be made to break up to two lines. The locus of poles of such degenerate cubics is the Hesse in H_2 , an $A^3 x^6$.^{**} To make these two lines coincide is three conditions and gives for the quartic an invariant, shown by Dr. Thomassen^{***} to be an A^{48} .

From the polar cubic curve can be obtained. The locus of poles of polar cubics having a double point is the Steinerian Σ , an $A^{12} x^{12}$. The cubic has also the invariants S and T , of degrees four and six, respectively, which give rise to the covariants of the quartic $S \equiv A^4 x^4$ [†] and $T \equiv A^6 x^6$ ^{††}. From the relations connecting the discriminant of the cubic with

* The notation $A^l x^m y^n$ is used to represent a covariant form of degree l in the coefficients of the quartic, m in x , and n in y .

** See note 1.

*** Am. Jour. Math., vol. XXXVIII, p. 249 (1916)

† See note 2.

†† See note 3.

the two invariant variables

$$T = 24x^3 + 1,$$

we may use it for calculating coefficients of the Steiner curve when that curve is necessary. It requires that the polar cubic to the curve is too complicated, giving rise to a set of twenty-four points in special polar curves. The

number of these points is determined by the fact that they are the common points of curve T ; therefore they are cusps of the Steiner curve. There are also twenty-one points whose polar cubics have two as multiple and therefore break up into a line and a curve; these are the double points of the Steiner curve. The line parts of these curves will hereafter be referred to simply as "twenty-one lines." Each of these lines is a straight line so that the four tangents at the intersection are on a straight line; in fact it is the straight line $z = 0$; i.e., the pole of the line is at the origin of the coordinate system. If one of these lines is called l as $x_0 = 0$ and the corresponding point is $(1, 0, 0)$, then the line is

$$v_0 = 0, v_1 = v_2 = 0, v_3 = 1.$$

The undulation is also a point of the Hessian, for its polar cubic is

$$hx_0^2 + 2b_0x_0x_1 + 2nx_0x_2 = x_0(hx_1 + 2b_0x_1 + 2nx_2),$$

a pair of lines. The double point of this is

as $(0, 0, 1)$, which therefore becomes the corresponding point on the Steinerian. Then

$$m = 0.$$

The polar cubic of $(0, 1, 0)$ is

$$a_1x_0^3 + 3hx_1^2x_0 + 3lx_0x_1^2 + 3b_0x_0x_1^2 + 3nx_0x_2^2.$$

This is made up of x_0 and a conic; its two double points are $(0, n, b_0\sqrt{-1})$ and $(0, n, -b_0\sqrt{-1})$; they are the two points of the Hessian corresponding to the double point of the Steinerian and are harmonic to the undulation point and the Steinerian point corresponding to the undulation considered as a Hessian point. The terms in H not containing x_0 are

$$x_1^4x_2^2 - cb_0^2 + x_1^2x_2^4 - 2cb_0n + x_2^6 - c^2v^2 = -cx_2^2(b_0x_1^2 + nx_2^2)^2.$$

Therefore x_0 is a triple tangent to the Hessian.

§ 3. The Discriminant of the Hessian.

The Hessian and Steinerian, as is well known, are not independent curves, but are brought into the same correspondence by the relations

$$\alpha x_1 \alpha_j^2 \alpha_i = 0, \quad i = 0, 1, 2,$$

where x is a point of the Steinerian and y a point of the Hessian. This says that the polar cubic of x has a double point y and the polar conic of y a double point x . The pairs of all such pairs x and y give rise to the Cayleyan, an $A'^2 g'^5$.

Since x and y cannot come together unless

$$\alpha x_i^3 = 0, \quad i = 0, 1, 2,$$

which is the condition that the quartic have a double point, there is a very convenient reference system for a study of these two curves in any characteristic. Let $(0, 1, 0)$ be a point of the Hessian, $(0, 0, 1)$ a corresponding point of the Steinerian, so that x_0 is a point of the Cayleyan. Then

$$x_2 x_1^2 x_i = 0, \quad i = 0, 1, 2,$$

$$v_1 = v_2 = 1 = C.$$

are each ordinary points.

The Hessian also has a double point at

$$c_1 = -2c_0.$$

The same property is satisfied by $(0, 0, 1)$, is also a double point, as

can be verified from the explicit forms. It is also a double point, as

is a double point, as $(0, 0, 1)$ is a double point, as

is a double point, as $(0, 0, 1)$ is a double point, as

$$4x_0^3 + 2c_0x_0x_1 + x_1^3$$

$$= 2x_0^3 + 2c_0x_0x_1 + c_0^2x_1^2.$$

if the same is conjugate of x_0 as to the first pair of lines is taken as x_1 , and that as to the second as x_2 , then

$$b_0 = c_0 = 0.$$

This shows that x_0 is one of the twenty seven lines. So the degree of the curve is 27. The same is one of the objects of the investigation.

With the above reference we have

$$J = x_1x_2^2 - b_0x_1^2$$

$$+ x_2^3 [x_0^2 (c_0^2 - 4b_0^2) + x_0x_1 - 2c_0x_1 + x_1^2] + c_0x_1^2 - c_0^2x_2^2$$

$$\Gamma = x_1 x_2^5 \cdot 2 b c^2 l^3$$

$$+ x_2^4 [x_0^2 \cdot 3 c^2 h^2 l^2 + 2 x_0 x_1 (-3 b c^2 a_2 h l + 3 b c^2 a_1 l^2)$$

$$+ x_1^2 (6 c^2 a_2^2 + 4 b^2 c l^3 + 12 b c^2 l^2)] + \dots \text{write } \dots \text{ in } x_2.$$

Therefore these two curves touch at $(0, 2, 1)$ along the line $x_1 = 0$ and

initially at $(0, 1, 0)$ and $q \neq 0$ in x_2 .

$$\Gamma = x_1^2 x_2^{10} + 3 c^2 l^6$$

$$+ x_2^9 [x_0^2 x_1^2 - 12 b c^4 h^2 l^5 + x_0 x_1^2 \cdot 3 c l^3 (-3 b c^2 a_2 l + 3 b c^2 a_1 l^2)$$

$$+ x_1^3 \cdot 4 b c^2 l^3 (6 c^2 a_2^2 - 12 b^2 c l^3 + 12 b c^2 l^2)] + \dots \text{write } \dots \text{ in } x_2.$$

Therefore Γ has a singularity at $(0, 0, 1)$, and symmetrically at

$(2, 1, 0)$, which is something more than merely a cusp. It may be

a tac-node.

These three cases make up the totality of ways in which the intersection may acquire a double point. The effect of its discriminant, Δ^{225} , must be added of the tree Γ is its attached to it.

§ 4. The discriminant of sf^4 .

The contour of sf^4 is, we know,

$$b_{\infty} = f = c - c.$$

The point of contact is given by

$$3c_1 m J_1 + (-b c_0 + b_0 c_1) J_2 = 0.$$

Since $(0, 0, 1)$ is taken as a ∞ of one of the points in which x_0 cuts the quartic, the points of contact will be on ax^4 of

$$x=0.$$

But this says that $(0, 0, 1)$ is a point of the Steinerian. Therefore the forty-eight intersections of ax^4 and sf^4 , which considered as a point curve is of order twelve, are the same as those of ax^4 and Σ . This can be substantiated by finding the

point equation of sf^4 . The line equation of ax^4 is $s^3 - 27t^2$.

Let s for ax^4 and sf^4 is $-12S + A^3$, and A^3 is $s^3 - 27t^2$.

Let A for ax^4 and sf^4 is $-12S + A^3$, and A^3 is $s^3 - 27t^2$.

$T + (ax^4 \cdot tx')^3 (tx'')^3 (x'x)(x''x)$. Therefore the equation of sf^4 is

$$\begin{aligned} & [-12S + A^3] \cdot (ax^4)^3 - 27 \Sigma [T + (ax^4 \cdot tx')^3 (tx'')^3 (x'x)(x''x)] \\ & = -27 \Sigma + (ax^4)^4 \cdot A'' x^5. \end{aligned}$$

x_0 will become a double line of s only if

$$b_0 = c_0 = m = 0,$$

If $c_1 = 0$, the quartic would have a double line.

all other conditions than this placing x_0 as a constant in the quadratic. The points in which x_0 meets the quadratic are $(0, 0, 1)$, $(0, \kappa, 1)$, $(0, \omega\kappa, 1)$, $(0, \omega^2\kappa, 1)$, where $\kappa^3 = -\frac{4c_1}{b}$ and $\omega^3 = 1$. The tangents to the quadratic at these four points are, respectively,

$$c_1 x_1 = 0,$$

$$3n\kappa x_0 - 3c_1 x_1 + 3c_2 x_2 = 0,$$

$$3n\omega\kappa x_0 - 3c_1 x_1 + 3c_2 \omega\kappa x_2 = 0,$$

$$3n\omega^2\kappa x_0 - 3c_1 x_1 + 3c_2 \omega^2\kappa x_2 = 0,$$

and they evidently have the common point $(c_1, 0, -n)$. Let

$$n = 0,$$

so that this common point becomes $(1, 0, 0)$. Since the coefficients equated to zero are b_2, n, b_3 , the point $(0, 1, 0)$ is a point of the Hessian with $(0, 0, 1)$ as its corresponding point.

Since the tangent to the cubic at $(0, 1, 0)$ is $x_1 = 0$, the point $(0, 1, 0)$ is a point of inflection. Similarly, the point $(0, 0, 1)$ is a point of inflection, and the point $(1, 0, 0)$ is a point of inflection. The point $(1, 0, 0)$ is a point of inflection, and the point $(0, 1, 0)$ is a point of inflection. The point $(0, 0, 1)$ is a point of inflection, and the point $(1, 0, 0)$ is a point of inflection.

The same distinction was made as to the four points

in which x_0 cuts the quartic by assigning the particular coordinates $(0, 1, 0)$ to one of them, then one of the two ty-one lines is a line of sf^4 , the quartic and the Steinerian touch four times on this line, the four tangents going thru the corresponding point. Also, since the tangent to the Hessian at $(0, 1, 0)$ is x_2 , the tangents at the four Hessian points corresponding to these Steinerian points are also on that corresponding point.

x_0 will also be a line of sf^4 if

$$b = b_2 = f = 0;$$

i.e., if x_0 is a stationary line of the quartic. But this leads to nothing new. Therefore sf^4 has a double line only when one of the two ty-one lines is on sf^4 , and its discriminant, t_0 , or A^{54} , expresses this condition.

§ 5. The discriminant of tf^6 .

The contravariant $(tf)^6$ is on the line x_0 if

$$b = b_2 = f = 0.$$

Then

$$(t f)^6 = 2 f_0^5 f_1 + 2 b_0 c_1 x^2 + \text{lower powers in } f_0.$$

x_0 becomes a double line either if $b_0 = 0$, which says that the quartic has a double point at $(c, 0)$ for x_0 as one of the tangents there, or if $c_1 = 0$, which says that the quartic is a straight line.

$(t f)^6$ is also on x_0 when

$$b_2 = f = c_1 = 0.$$

Then

$$(t f)^6 = 2 f_0^5 f_1 + b c x + 2 f_0^5 f_2 + b c x + \text{lower powers in } f_0.$$

For x_0 to be a double line requires either $b=0$ or $c=0$, which repeats the undulation condition, or $m=n=0$, which is the condition on the quartic under which the Hessian has two double points and for which the invariant of undulation degree vanishes. Then one of the two tyrosolines is a line of $(t f)^6$:

$(t f)^6$ also has x_0 as a line when

$$b = f = c = 0,$$

but this leads to $\text{not } f$ in $\mathcal{F}$. Therefore the discriminant

of $(t\mathfrak{f})^6$, an A^{220} , must be made up of the discriminant of the quartic, an A^{27} , the modulation in coefficients, an A^{10} , and the unknown invariant.

§ 6. The Twenty-one Lines.

The twenty-one lines* are given by an $A^i x^{21}$, where i is as yet unknown. That it can be determined is due to the fact that these lines consist of the common tangents of four conics, all of whose coefficients are known. They are, the tangents to $A'^{12}f^{18}$, and to $A'^{15}f^{24}$, and to the two of f touching the quartic so that three of the tangents at the triple section are in a point.** The common lines of these conics are:***

* A curve on these lines is

$$2(s\mathfrak{f})^3(s\alpha)(s'\mathfrak{f})^3(s'\alpha)(s''\mathfrak{f})(s''\alpha)(s'''\mathfrak{f})^3(s'''\alpha) - 3s\mathfrak{f}^4(s'\mathfrak{f})^3s'\alpha(s''\mathfrak{f})^3s''\alpha(s'''\mathfrak{f})^2(s'''\alpha)^2 \\ + \frac{1}{3}s\mathfrak{f}^4(s'\mathfrak{f})^4(s''\mathfrak{f})^4(s'''\mathfrak{f})^4.$$

** Am. Jour. Math., vol. XXXIX, p. 227 (1917).

*** Let $v_2 = v_1 = 0$, so that x_0 is a double point. Let

such invariants as express the fact that x is b.s. or t is del-
 ete. The q such invariants is a q such invariant of x or t is a q such invariant, for if the q such invariant is a q such invariant it di-
 vides x at least once for each of x and t . It is

$$A^{558} x^{432} = (A^i x^{21})^{16} \cdot (A^{14} x^{24})^2 \cdot A^{10} x^{48} \cdot (A^{27})^k$$

$$\therefore 558 = 192 + 108 + 270.$$

$$192 + 270 = 462.$$

It is evident that k shows that i must be a multiple of 18 and
 can be satisfied only when $i=18$. Therefore the twenty-one
 lines are given by $A^{18} x^{21}$.

§ 7. The Invariant of Unknown Degree.

The degree of the form giving the twenty-one lines being
 known, we can force one of them to lie on either sg^4 or tg^6 .
 The condition so obtained will contain the undulation
 condition, however, for the tangent at an undulation is
 a special case of one of the twenty-one lines and is a line of
 both sg and tg . Suppose the $A^{18} x^{21}$ factored into its components

A^{15} is, the coordinates of each of these substituted in tf_1^6 , the
 result of the substitution is $A^{15} = A^{15}$, and the
 coefficients of the $A^{15} x^{21}$ substituted for A^{15} is the
 result of these coordinates. The result is

$$A^{15 \cdot 4 + 21 \cdot 21} = A^{114} = A^{60} \cdot A^{54}.$$

It is clear that tf_1^6 is a degree of the required invariant. For the invariant

tf_1^6 is a degree of the required invariant. For the invariant

$$A^{18 \cdot 6 + 21 \cdot 3} = A^{171}$$

from which the invariance condition must

divide out at least 111. Therefore the invariant in question

may be either of degree 111 or of degree 51. The first of these

will not fit into the diagram of the invariant, but the

$$A^{225} = A^{21} \cdot (A^{48})^2 \cdot (A^{51})^2,$$

while for the invariant of tf_1^6 we have

$$A^{225} = A^{21} \cdot (A^{60})^2 \cdot A^{51}.$$

Therefore we have a invariant of degree 51, the result of which

expresses the invariant of the invariant as a function of the

tf_1^6 as a degree of the invariant, and the invariant of the invariant is a

of tf_1^6 . A further meaning will be seen later.

§ 8. Certain ...

The degree of the form giving the twenty-one lines being known, that for the twenty-one corresponding points can be calculated from the fact that each point is a base point of the corresponding line as to both sg^4 and g^6 . Suppose, as before, the $A^{18}x^{21}$ factored into its component lines, the coordinates substituted for x in sg^4 and the symmetric functions of these coordinates in the product of the resulting terms replaced by the coefficients of $A^{18}x^{21}$. The result gives the twenty-one points together with such invariants as express that the polar point lies on the line, i.e., that one of the lines be a double line of sg^4 .

$$A^{18 \cdot 3 + 2 \cdot 21} g^{21} = A^{54} \cdot A^{42} g^{21}.$$

Similarly for sg^6 , we obtain

$$A^{18 \cdot 5 + 3 \cdot 21} g^{21} = A^{60} \cdot A^{51} \cdot A^{42} g^{21}.$$

The representation of the points as a variety is $A^{42}x^{21}$. *

* This degree fits in with the form $A^{36}g^{18}$ for the line equation of Σ . For the cusps are counted for three times in each of S, T, X as in $A^{48}g^{24}$. Therefore the Plücker formula

$$A^{12 \cdot 22} g^{12 \cdot 11} = A^{264} g^{132} = A^{42} g^{21 \cdot 2} \cdot (A^{48} g^{24})^3 \cdot A^{36} g^{18}.$$

one of them is on $(ax)^4$ (and therefore is its own reciprocal).

Then, as before, consider $A^{42}g^{21}A^{10}$ to its reciprocal

points, the coordinates will be x^4 , and x^4 is replaced in the product of the result-

ing terms, we obtain $A^{42+4+21} = A^{189}$. Out of this the

condition for an undulation, when the point is on the line part of its parabolic, must divide out at least once.

If it is so more than once, we have left A^{129} to express that the point is on the cubic part of its parabolic. It

suffices that x is degree 129 and x^4 is degree 129.

Since the condition is not 129, we had before that the point is $(1,0,0)$ and its corresponding line x_0 if

$$b_0 = m = n = c_0 = c.$$

To put the point on the quartic requires that

$$x = 0.$$

The undulation case is excluded because no choice of coordinates for the point and line is possible that they are not incident. Recall that $(0,1,0)$ and $(0,0,1)$ are x_0 .

Let us take x_1 as the new roots of the equation of $1, 0, 1, 20x_1 + t$

$$q = 1 - 0.$$

If q is sf^4 the coefficients of $J_1^4, J_2^4, J_1^3 J_2, J_1 J_2^3$ are $1, 5, 10, 10, 1$ so that $1, 0, 1, 20x_1 + t$ is a point of sf^4 .^{*} It is known that the double lines of sf^4 are given by $A^{16}x^{28}$; therefore the double points of sf^4 are given by $A^{32}f^{28}$. If q is sf^4 , then sf^4 is a double point, so then $(sf^4)^4$ has that same double point with the same tangents. Then by the method of considering the $A^{32}f^{28}$ factored into its component parts, we have as the condition that q is a double point to sf^4

$$A^{32 \cdot 4 + 28} = A^{156} = A^{129} \cdot A^{27}.$$

This establishes the degree of the A^{129} .

The totality of polar cubics of the twenty-one points given by the $A^{42}f^{21}$ is given by $A^{63}x^{63}$. Suffice it is

* An examination of the coefficients of sf^4 shows that the condition is not a double point of it.

factored into its component cubics $(p_i x)^3$, and then each of these cubics factored into its component line and conic $(y_i x) \cdot (\delta_i x)^2$. Then the condition that the line touch the conic is of second degree in both y_i and δ_i , therefore of second degree in p_i . Therefore the invariant expressing the condition that there be a polar cubic made up of a conic and a line touching it is an $A^{6 \cdot 3 \cdot 2} = A^{126}$. This result can be substantiated in other ways. If, as usual, we take the line as x_0 and the corresponding point as $(0, 0, 0)$, so that

$$b_0 = m = n = c_0 = 0,$$

then the conic will touch x_0 at $(0, 1, 0)$ if

$$ah = b = 0.$$

Then S goes thru $(0, 1, 0)$ while T has a double p. there, so that Σ has ~~two~~^{two} double points and two cusps coming together and acquires a triple point, since x_0^9 is the highest power of x_0 recurring in its equation. Since two of the intersections of S and T have come together, the A^{126} must be a factor of the tact-invariant of these two curves. We have seen (p. 8, 10) that in case of the intersection of either

the A^{48} or the A^{51} that S and T touch at two distinct points, and these three cases seem to be the only ones in which two intersections of these curves come together. Their tact-invariant is found from the formula given by Salmon to be of degree $4 \cdot 6 (6+8-3) + 6 \cdot 4 (4+12-3) = 576$. Then

$$A^{576} = (A^{48})^2 \cdot (A^{51})^2 \cdot (A^{126})^3.$$

Also inspection of their coefficients shows that both H and S touch x_0 at $(0, 1, 0)$. As will be seen later, two intersections of S and H can come together only when two of S and T do. When the A^{51} vanishes, H has two double points on S ; when the A^{48} vanishes, H has a double point which is also a double point of S . The tact-invariant of S and H is of degree $4 \cdot 6 (6+8-3) + 3 \cdot 4 (4+12-3) = 420$.

$$A^{420} = (A^{48})^4 \cdot (A^{51})^2 \cdot A^{126}.$$

The twenty-one polar cubics being given by an $A^{63}x^{63}$, while the twenty-one line parts are given by $A^{12}x^{12}$, the twenty-one conic parts are given by an $A^{45}x^{45}$. Since the discriminant of a conic is of third degree in its coefficients, the condition that one of these conics break up

is an A^{45} . But then the polar cubic consists of three lines, any two of which may be considered as the cubic part, so that this degree is probably divided by three. Therefore the condition that there be a polar cubic made up of three lines is an A^{45} . Then three of the double points of the Steinerian have come together, giving it a triple point.*

* Professor Morley has pointed out to me that the degrees of these last two invariants can be substantiated by the work of Caporali. See *Memorie di Geometria*, p. 171, on a web of four curves. If n curves of a web are of a particular kind, then the invariant expressing that there be a curve of that kind is a function of degree n in the coefficients of the curve upon which the invariant is a function. From his results the invariants expressing that in a set of four there be a curve of a particular kind, or a curve with two or more like tangent points, are of degrees 15 and 42, respectively, as recorded in the table of values.

It is supposed that each invariant is linear in the coefficients of the quartic. Therefore the invariants are of degree 45 and 126 in the coefficients of the quartic.

§ 9 The Eliminant of S, T, H .

Under the reference scheme here

$$b_2 = f = n = 0,$$

the coefficient of the highest term in x_2 in both S and T is a power of $c, b + v^2$. Then both coefficients vanish if $c = n = 0$; it is, as we have seen, also the vanishing of f on A^5 . Then for the general quartic we obtain $(0, 0, 1)$ as one of the c, b, v points of S and T if we make

$$b = n = 0.$$

But then the coefficient of x_1^4 vanishes in S . Therefore to be sure that four cusps of the intersection correspond to one vertex of the tetrahedron of the Hessian then each of its vertices is on S .

It is now what the plane curve of a point on S is reduced to the curve of the intersection of the plane with its Hessian is the product of the factors ϕ and ψ of $\phi = 0$ and $\psi = 0$ is $\phi\psi$. If the curve has a double point there is only one such ϕ and ψ made up

of the tangents at the double point and a line through the three remaining flexes flex. i.e., the surface cut here will be tangent to the curve tangent counted twice and the line will be tangent to the curve to the surface remaining flex.

Here the polar cubic of $(0, 0, 1)$ is

$$a_2 x_0^3 + 3g x_0^2 x_2 + 3c_0 x_0 x_2^2 + 3c_1 x_1 x_2^2 + c_2 x_2^3,$$

and its Hessian is

$$c_1^2 x_2^2 (a_2 x_0 + g x_2).$$

But x_2 is the tangent to H at $(0, 1, 0)$ and $a_2 x_0 + g x_2$ is the tangent to S at the same point. Therefore the polar cubic of a cusp of the surface has as its cusp the point and its flex the point tangent to H and S , respectively, at the corresponding point. The cubic curve is thus tangent to the surface.

$$\frac{1}{a_2^2} \left[(a_2 x_0 + g x_2)^3 + x_2^2 \{ 3a_2(a_2 c_0 - g^2) x_0 + 3a_2^2 c_1 x_1 + (c a_2^2 - g^3) x_2 \} \right]$$

which also shows that the tangent at the flex is the surface in brackets

The polar cubic of $(0, 1, 0)$ is

$$a_1 x_0^3 + 3b_0 x_0^2 x_1 + 3b_0 x_0 x_1^2 + b_1 x_1^3 + c_1 x_2,$$

and its Hessian is

$$c_1 x_2 \left[x_0^2 (a_1 b_0 - h^2) + x_0 x_1 (b a_1 - b_0 h) + x_1^2 (h b - b_0^2) \right].$$

Then the tangent to H at $(0, 1, 0)$ passes thru three flexes of the cubic, the three flex tangents pass thru the corresponding Steinerian point and the binary Hessian of these three tangents picks up the six remaining flexes.

The coefficient of x_2^2 in H is $-c_1^2 g$. Since we can not have $c_1 = 0$ without bringing on the vanishing of the $A^{(5)}$ and worse, let us take

$$g = 0.$$

Then we have made S, T, H have a common point at $(0, 0, 1)$.

The polar cubic of $(0, 0, 1)$ becomes

$$a_2 x_0^3 + x_2^2 (3 c_0 x_0 + 3 c_1 x_1 + c_2 x_2).$$

If we let the intersection of the stationary tangent lines be $(0, 1, 0, 0, 1, 2, 2)$

$$c_0 = 0.$$

(But then, since we have $g = n = c_0 = 0$, the Steinerian point corresponding to $(0, 0, 1)$ as a Hessian point is $(1, 0, 0)$, which also becomes a cusp on the Steinerian. So to sum up, we have: 1) the cusp e_0 on the Steinerian corresponds to $e_2, 0, 0$

§ 10. The Eliminant of the Polar Cubic, Conic, and Line.

Suppose we ask that the polar cubic, conic, and line of $(0, 1, 0)$, which is not on the quartic, have a common point $(0, 0, 1)$.

Then

$$b_2 = f - c_1 - 0.$$

The symmetry of this condition shows that the relation ~~bet~~ between $(0, 1, 2)$ and $(0, 0, 1)$ is a natural one. Also, x_0 is a line of $\mathcal{H}^6$. Therefore we might define $\mathcal{H}^6$ as the locus of the joins of pairs of points so related that the polar curves of each pass thru the other, the pair of points being the common harmonic pair of one of the harmonic pairs of points in which their join cuts the quartic.

In general the skin in a t of the placenta is x , which

are an Ax^3y , $\sim Ax^2y^2$, and $\sim Ax^4$, is an $A''x^{26}$. But if this
 x^6 divides twice, i.e. $\sim y^2 \sim A^9x^9$. Def. exactly y^2 is the
 of x is "little" $\sim y^2$, let us say. The other x^6 is a "little" $\sim y^2$
 Ax^4 , its polar is Ax^2y^2 , and its polar is Ax^2y^4 . It is an $A''x^{13}$,
 for x^6 which the cubic divides out twice. The x^6 is a "little", a
 A^9x^9 , gives x^6 a "little" stationary line. This for the A^9x^{13} is y^2
 the y^2 is a "little" $\sim y^2$, so the x^6 is of it $\sim y^2$ is a "little" $\sim y^2$
 the x^6 is a "little" $\sim y^2$, so the x^6 is of it $\sim y^2$ is a "little" $\sim y^2$
 the x^6 is a "little" $\sim y^2$, so the x^6 is of it $\sim y^2$ is a "little" $\sim y^2$

Salmon gives as the equation of the nine stationary
 lines of the cubic

$$S \circ H^2 - H^3 - 10D,$$

where S is the invariant of degree four, H is the cubic itself,
 H is its Hessian, and D is a certain A^3x^6 . For a polar conic
 of the quartic the first three possessive planes are $\sim y^2$ to
 the invariant S , the quartic itself H , and its Hessian, and D
 is not so easy to understand. So we will substitute for it
 another A^3x^6 , expressible in terms of S , H , and D , which
 gives the locus of points whose polar conics as to the cubic

are on their polar lines as to the Hessian. The result is easily shown that

$$A^9 x^{18} = 8 S d^2 H - H^3 + 9 d d O',$$

where S is the invariant S of the quartic, d the quartic, H its Hessian, and O' the locus of those points whose polar lines as to the quartic are on their polar lines as to the Hessian.

The form of the $A^9 x^{18}$ shows that all of its "intersections with O' " are used up at the flexes of the quartic. These twenty-four points are also part of its intersections with H , if it there are eighty points, lying on O' . Now we saw that $0, 1, 0$ is a point of the $A^9 x^{18}$ if

$$b_2 = f = c = 0.$$

To make it a point of H also requires that

$$a = 0. \quad *$$

But there are only forty-two such points, for they are the points where H, O' and the Cayley curve all to meet. ^{**} Hence the intersections of H and O' must come to set eleven pairs, ²

* Or, symmetrically, $0, 0, 1$ is a point of H if $a = 0$.

** Proc. Nat. Ac. Sci., vol. 3, p. 449 (1917)

indeed, it is readily shown that the two curves touch at their common points.

It is also easily shown that the forty-two inflexions of the Steinerian are at the points corresponding to these Hessian points.

§ 11. The Polar Conic of Two Points.

The invariant of the polar conics of two given points is the discriminant of the polar conic of two points in the plane of x, y, z . The polar conic of x, y, z as to the quadratic is $(dx, dy, dz)^2$. If y point on the line xy may be represented parametrically as $x + \lambda y$. This line meets the conic in two points whose parameters λ are given by

$$(dx, dy, dz)(dx + \lambda dy)^2 = dx^3 + 2\lambda(dx)(dy)^2 + \lambda^2(dx)(dy)^3 = 0.$$

If we ask that these points become coincident, we require that the coefficient to λ^2 shall vanish. This is the case if $(dx, dy, dz) = 0$, which is the condition for the line xy to be a tangent to the conic. If we ask that the two points become coincident, we require that the coefficient to λ shall vanish. This is the case if $(dx, dy, dz) = 0$, which is the condition for the line xy to be a tangent to the conic.

of the first series

$\frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

$$v_2 = f = e_1 = 0.$$

2. 2 p v a v c i e , t e t r o f s t o s

$$y_2'' + 2yx_0x_1 + 2yx_0x_2 = x_0(y_0 + 2yx_1 + 2yx_2).$$

So ϕ is a $\mathbb{Z}_p$ -homomorphism to G is the

$$A = C \quad , \quad r = 0$$

Let τ is axis $(0, 1, 0)$ and if we plug it into ρ , it becomes 6.

[illegible]

ex. 1: stand x at the end of the street, then x is a
 point of the axis x if a point in the interval $(0, 1)$.
 the point is $(0, 1)$ and $(0, 1)$ is

$$lx_0^2 + 2 \quad u_0 x_1 + 2 \quad u_0 x_2 + u_2 x_1 + 2 \int x_1 x_2 + c_1 x_{12}$$

is not a new is ... 0, 1, 0, 1

$$v_0 = f = 0,$$

be y , x and z respectively at the origin of the curve of the t axis. So as x and z are x and z of the t axis at the origin of the t axis, t , as we have seen, the forty-two flexions of the H curve and one forty-two flexions of the H curve are the angles α and β .

Now if a point t , let t be the polar coordinate of $(0, 1, 0) = a$ $(0, 1, 1)$ be the square of a line. If we take $(0, 1, 1)$ as a point of this line, then

$$l = m = n = 0,$$

$$b_2 c_1 - f^2 = 0.$$

But then we find that the coefficients of x_1^4 and x_2^4 in the equation are therefore the pair of points whose polar coordinate is a repeated line in $\mathcal{S}$. But we already have a correspondence of the points of $\mathcal{S}$, and the polar coordinate of a point a , $\mathcal{S}$ is a line of three lines, the intersections of which are in $\mathcal{S}$. With the above reference we have

$$v_{4,0} = (x_1 v_0 - v^2)(-x_2 c_0 + f^2).$$

$(1, 0, 0)$ was any point of the repeated line. Let us take it as one of the intersection points in S by noting

$$a, b_0, v^2 = 0.$$

Then the projection of $(1, 0, 0)$ becomes

$$x_1, x_0 x_1 (b a_1 f - a_1 v_2^2 - v_0^2 f h) + x_0 x_2 (v a_1 c_1 - a_1 v_2 f - v_0 c_1 v) \\ + x_1^2 (v_1 v - b_0^2 f - v_2^2 v) + x_1 x_2 (b c_1 v - v_0^2 c_1 - v_2 v) = 0.$$

The double point is $(1, 0, 0)$, $(0, 0, 1)$ and

$$f = (v c_1 v - v_0^2 c_1 - v_2 f v, -b a_1 c_1 + a_1 v_2 f + b_0 c_1 v, v a_1 f - a_1 v_2^2 - b_0 f h).$$

Then the tangent line of $(0, 1, 0)$ and $(1, 0, 0)$ is $a_1 x_0^2 + 2 f x_0 x_1 + v_0 x_1^2$,

that of $(0, 1, 0)$ and $(0, 0, 1)$ is $b_2 x_1^2 + 2 f x_1 x_2 + c_1 x_2^2$, and that of $(0, 1, 0)$

and $(1, 0, 0)$ is x_1^2 . The equation of the plane is of a point in S and a line

is the projection of the intersection line is the equation of the plane. *

* We might expect this sort of three-to-one correspondence from the following considerations: The line equation of x_0, x_1, x_2 is an $A^2 x^2 y^2 z^2$. If the curve is the square of a line the line equation vanishes identically. Equating each coefficient to zero, we have six equations from which we can eliminate the six functions $f_0^2, f_1^2, f_2^2, f_0 f_1, f_1 f_2, f_2 f_0$, leaving $A^2 x^2$. It is not true.

If we also ask that it repeated be zero $(0, 1, 0)$ we

$$b_2 = g = 0.$$

Then $(0, 1, 0)$ is one of the twenty four points of S and H .
Therefore the polar conic of a SH point and its corresponding T point is the square of the line SH point T line
tangent to H line.

If, finally, we ask that the polar conic of $(1, 1, 0)$ and
 $(0, 0, 1)$ be x^2 , we have

$$m = n = b_2 = f = c_1 = 0.$$

But then the A^5 vanishes. Therefore we have a new definition for the A^5 ; it is the invariant expressing the condition that there be a pair of points such that their polar conic as to the quartic is the square of the line joining them.*

§12. Salmon's Connex.

Salmon has shown that with any plane curve of n there

* See note 4.

is associated an $A^3 x^{2(n-2)} y^{n-2}$, which when x is a point on the curve picks out the remaining intersections of the tangent at x with the curve. This connex is expressible in terms of polars of the Hessian of the curve itself and the Hessians of the polar curves of x as to $(xy)^n$. In case of the quartic the connex is an

$$A^3 x^4 y^2 \equiv 15(hx)^4(hy)^2 - 9(h, x)(h, y)^2,$$

where $(hy)^6$ is the Hessian of the quartic and $(h, y)^3$ is the Hessian of the polar cubic of x . Explicitly

$$\begin{aligned} A^3 x^4 y^2 = \sum^3 \left[x_0^4 \{ y_0^2 \cdot 6(a_0 h - a_1^2 - a_1^2 g - a_2^2 h + 2a_1 a_2 l) \right. \\ + y_0 f_1 \cdot 4(ab_0 g + ahn - 2alm - a_2^2 b_0 - a_1^2 n \\ + 2a_1 a_2 m - a_1 gh + a_1 l^2) \\ + f_0 f_2 \cdot 4(ac_0 h + agm - 2aln - a_1^2 c_0 + 2a_1 a_2 n \\ - a_2^2 m - a_2 gh + a_2 l^2) \\ + f_1^2 (abg - ba_2^2 + 2b_0 n - 2ab_2 l + afh - a_1 n^2 \\ + 2a_1 a_2 b_2 - a_1^2 f - a_1 b_0 g - a_1 n v + 2a_1 l n, \\ + f_1 f_2 (ab_0 c_0 + 2ab_2 g + 2ac_1 h - 4afl - amn - 2a_1^2 c_1 \\ + 2a_2^2 b_2 + 4a_1 a_2 g - a_1 c_0 h - a_2 b_0 g \\ - a_1 g n - a_2 h v + 2a_1 l m + 2a_2 l n, \end{aligned}$$

$$\begin{aligned}
& + f_2^2 (aeh - ca_1^2 + ac_0 n - 2ac_1 l + afg - a v^2 + 2a_1 a_2 c_1 \\
& \quad - a_2^2 f - a_2 c_0 h - a_2 g n + 2a_2 l n), \\
& + x_0^3 x_1 (f_0^2 \cdot 8 (ab_0 g + ah n - 2al n - a_2^2 b_0 - a_1^2 v + 2a_1 a_2 n \\
& \quad - a_1 gh + a_1 l^2)) \\
& + f_0 f_1 \cdot 2 (abg - ba_2^2 + 7ab_0 n - 2ab_2 l + afh - 7a_1 n^2 \\
& \quad + 2a_1 a_2 b_2 - a_1^2 f + 5a_1 b_0 g - 12a_2 b_0 l \\
& \quad - 7a_1 h n + 12a_2 h n + 2a_1 l n - 6gh^2 \\
& \quad + 6nl^2) \\
& + f_0 f_2 (7ab_0 c_0 + 2ab_2 g + 2ac_1 h - 4afl - 7a_1 n n - 2a_1^2 c_1 \\
& \quad - 2a_2^2 b_2 + 4a_1 a_2 f - 7a_1 c_0 h - 1a_2 b_0 g + 17a_1 g n \\
& \quad + 17a_2 h n - 10a_1 l n - 10a_2 l n - 12ghl \\
& \quad + 12l^3) \\
& + f_1^2 \cdot 3 (abn + ab_0 f + ba_1 g - 2ab_2 n - 2ba_2 l \\
& \quad + 2a_2 b_2 h - a_1 jh - b_0 gh - h^2 n + 2kl n) \\
& + f_1 f_2 (abc_0 + 5ab_0 c_1 - ba_2 g + 3ab_2 n - 9af n \\
& \quad + 2a_1 b_0 c_0 + 1a_1 b_2 g - 5a_1 c_1 h - 4a_2 b_0 n \\
& \quad - 10a_2 b_2 l + 11a_2 l n - 3c_0 v^2 - 2a_1 l \\
& \quad - 5b_0 g n - 2a_1 n v + 1a_2 j^2 - 3: n \\
& \quad + 3 \quad , + l^2 \quad)
\end{aligned}$$

$$\begin{aligned}
& + f_2^2 (2ab_0c_0 + 2b_2c_0 - 2ca_1v - 3ac_1v - 2a_2b_0c_0 \\
& - a_2b_2q + 5a_2c_1v + 4a_1c_0v - 2a_1c_1l \\
& + 4a_1fq - 4a_2ql - 3c_0l - 4a_1v^2 + 2a_2v \\
& - 3qlv + 3v^2v); \\
& + x_0^3 x_2 (f_0^2 (8(ac_0v + 2q_1v - 2a_1v - a_1^2c_0 + 2a_1a_2v - a_2v \\
& - a_2q_1v + a_2l^2) \\
& + f_0f_1 (7ab_0c_0 + 2ab_2q + 2ac_1v + 4af_1v - 7a_1v \\
& - 2a_1^2c_1 - 2a_2^2v_2 + 4a_1a_2f_1 - 7a_1c_0v \\
& - 7a_2v_0q + 11a_1q_1v + 17a_2v_1v - 10a_1c_1v \\
& - 10a_2l_1v - 12a_2l + 12l^3) \\
& + f_0f_2 (2(ac_1v - ca_1^2 + 7ac_0v - 2ac_1l + af_1q - 7a_1v^2 \\
& + 2a_1a_2c_1 - a_2^2f_1 + 5a_2c_0v - 12a_1c_0l \\
& + 12a_1q_1v - 7a_2q_1v + 2a_2v_1v - 3q_1^2v \\
& + 3ql^2) \\
& + f_1^2 (2ab_0c_0 + ab_0c_1 - 2ab_2q - 3ab_2v - 2a_1v_0c_0 \\
& + 5a_1b_2q - a_1c_1v + 4a_2v_0v - 2a_2b_2v \\
& + 4a_2f_1v - 4a_1ql - 3v_0ql + 2a_1v_1v - 4a_2v_1^2 \\
& - 3v_1^2v + 3v_1^2v); \\
& + f_1f_2 (ac_0l + 5ab_2c_0 - ca_1v + 3ac_1v - 9af_1v
\end{aligned}$$

$$\begin{aligned}
& + 2a_2 b_0 c_0 - 5a_2 b_2 g + 7a_2 c_1 h - 4a_1 c_0 n \\
& - 10a_1 c_1 l + 11a_1 f g - 3b_0 g^2 - 2a_2 f l \\
& - 3c_0 k + 4a_1 n^2 - 2a_2 n v - 3g h v \\
& + 3g l v + 3b^2 n v) \\
& + f_2^2 \cdot 3(a c_0 v + a c_0 f + c a_2 v - 2a c_1 v - 2c a_1 v + 2a_1 c_0 g \\
& - 2a_2 f g - c_0 g v - g^2 v + 2g l v), \\
& + x_0^2 x_1^2 \{ f_0^2 \cdot 3(a b g - 3a_2^2 + 3a v_0 v - 2a b_2 l + a f h - 3a_1 v^2 \\
& + 2a_1 a_2 b_2 - a_1^2 f + a_1 b_0 g - 4a_2 b_0 l \\
& - 3a_1 n v + 4a_2 v v + 2a_1 l v - 2g n^2 + 2l^2) \\
& + f_1^2 \cdot 6(a v v + a v_0 f + a_1 g - 2a v_2 v - 2a v_2 l \\
& + 2a_2 v_2 v + a_1 v_0 v - v_1 v - v_0 v \\
& - a_1 v^2 - 4v_0 v^2 - 5v^2 v + 10l v) \\
& + v_0 f v \cdot 3(a c_0 + v_0 c_1 - 2a_2 f - a v v - a f v + 2a_1 c_0 \\
& + 3a_1 v_2 g - 2a_1 c_1 v + 4a_2 v v - 2a_2 v_2 v \\
& + 3a_2 v v - 3c_0 v^2 - 2a_1 f l - 7b_0 f l \\
& - 2a_1 v v - a_2 v^2 + 5g l v - k l v \\
& + 3l^2 v) \\
& + f_1^2 \cdot 3(2b f - 2b^2 + 3(a_1 v - 2a_2 v + g h - 3v^2 \\
& + 2a_2 v_2 + a_1 b_0 f - v_0^2 g - 4a_1 v v)
\end{aligned}$$

$$-3b_0h + 4b_2hl + 2b_0v - 2fh^2 + 2v^2 - v^2)$$

$$+ f_1 f_2 (3(abc_1 + ba_1c_0 - ab_2f - la_2v - bcl + 2a_1b_0c_1 + 3a_2b_0f - b_0c_0h + 4a_1v - 2a_2v_2v + 3v_2h) - 3c_1v^2 - 1a_1f - 2v_1g - 2v_0l - b_2v^2 + 5ffl - v - v + 6l - v^2)$$

$$+ f_2^2 (abc - 2b_2c_1 - ba_2c_0 + 2ca_1v_0 - 3ev^2 + 4a_1v_2c_0 + 4a_2v_0c_1 - 2a_1c_1v - 3b_2c_0l - 3a_2f - 3v_2al + 3c_0v - 3c_1vl + 5fah - 3h^2 - 3g - 3v^2 - 3v_1v^2 + 12l - v - f)$$

$$+ x_0^2 x_1 x_2 (f_0^2 \cdot 3(3ab_0c_0 + 2av_2f + 2ac_1v - 4a - 3a - 2a_1^2c_1 - 2a_2^2v - 2 + 4a_1a_2f - 3a_1c_0v - 3a_2v_0g + 5v_1f - 5a_2v - 2a_1 - 2a_2v - 4v - 4v + 4v))$$

$$+ x_0 f_1 \cdot 3(2a_1c_0 + 3a_1c_1 - 2a_2f + 2a_1v - 3a_1f + 2a_1c_0 + 5v_1 - 3a_1c_1 + 4a_2v - 3a_2v_0 - 4v_1v - 1v_0^2 - 2v_1v - 1v - 2v_1v - 2v_1^2 + 7v + 1v)$$

$$\begin{aligned}
& + f_0 f_2 \cdot 3(a_1 b_0 + 3 a_1 b_2 c_0 - c a_1 h + a c_1 - 5 a_1 f v \\
& + 5 a_2 b_0 c_0 - 3 a_2 b_2 q + 5 a_2 c_1 h + 4 a_1 c_0 v \\
& - 6 a_1 c_1 h + 7 a_1 f q - 7 b_0 q^2 - 2 a_2 f h \\
& - 13 c_0 h k - 4 a_1 v^2 - 6 a_2 v v + q g h v \\
& + q h v + 10 b^2 v)
\end{aligned}$$

$$\begin{aligned}
& + f_1^2 \cdot 3(a b c_1 + 2 b a_1 c_0 - a b_2 f - 3 b g h + 4 a_2 b_0 f \\
& - 2 c_0 b h - 4 a_2 v_2 v + 5 b_2 q v - c_1 v^2 \\
& - 2 a_1 f v - 2 c_0 q v - (4 h k + 4 h v^2))
\end{aligned}$$

$$\begin{aligned}
& + f_1 f_2 (a b c + 8 a b_2 c_1 + 2 b a_2 c_0 + 2 c a_1 v_0 - 9 a f^2 - 3 b g^2 \\
& - 3 c h^2 + 10 a_1 b_2 c_0 + 10 a_2 b_0 c_1 - 12 a_1 c_1 v \\
& - 12 a_2 b_2 v - 5 a_1 f v - 5 a_2 f v \\
& - 18 b_0 q v - 12 b_2 c h - 13 c_0 v v \\
& - 12 c_1 h + 30 f g h + 6 g v^2 + 5 h v^2 \\
& + 24 h v v)
\end{aligned}$$

$$\begin{aligned}
& + f_2^2 \cdot 3(a c b_2 + 2 c a_2 v_0 - a c_1 f - 3 c_1 h + 4 a_1 c_0 f \\
& - 2 c_0 c_0 q - 4 a_1 c_1 v - b_2 q^2 + 5 c_1 q h \\
& - 2 a_2 f v - 2 c_0 v f g h + 4 h v^2)
\end{aligned}$$

This can be easily considered either as a vector field or as a differential equation depending on whether x or y is a coordinate or not. If it is a coordinate,

as a vector field, the corresponding vector field lies
 at t

$$x - x_1 = x_2 = 0,$$

so that $(1, 0, 0)$ is a critical point, and let us assume that
 point t

$$A^{3/4} x^2 = x_0^2 x_1^2 \cdot 3(-2gh^2 + 2cl^2)$$

$$+ x_0^2 x_1 x_2 \cdot 3(-4gh + 4cl^2)$$

$$+ x_0^2 x_2^2 \cdot 3(-2gh^2 + 2cl^2)$$

$$+ x_0 x_1^3 \cdot 3(-2gh - 2cl^2 + 2ghn + 2cl^2n)$$

$$+ x_0 x_1^2 x_2 \cdot 3(-c_0 h^2 - 2b_0 gl - ghn + 4l^2n)$$

$$+ x_0 x_1 x_2^2 \cdot 3(-b_0 g^2 - 2c_0 hl - ghv + 4l^2v)$$

$$+ x_0 x_2^3 \cdot 3(-c_0 gh - g^2n + 2glv)$$

$$+ x_1^4 \cdot 3gh - 4bl^2 - b_0^2 g - b_0 hn + 2b_0 l n)$$

$$+ x_1^3 x_2 (-b_0 c_0 v + b_2 gh - 4b_0 g n + 2b_0 l v - 4b_2 l^2 - 3hn + 3n^2 + 3l^2 v^2)$$

$$+ x_1^2 x_2^2 (-3b_0 g n - 3c_0 h n + 3gh - 3l^2 - 3g n^2 - 3n v^2 + 12l n n)$$

$$+ x_1 x_2^3 (-b_0 c_0 g + c_1 gh - 4c_0 n v + 2c_0 l v - 4c_1 l^2 - 3g n v + 3l^2 v^2)$$

$$+ x_2^4 (cgh - 4cl^2 - c_0^2 h - c_0 g n + 2c_0 l v).$$

Obviously this curve has a double point at $(1, 0, 0)$ in the same tangent to the quadric. Therefore six of the intersection points of the two curves are used up here. If needed the others let us take $(0, 1, 0)$ or x_1^4 , so that

$$b = 0.$$

Then in order that $(0, 1, 0)$ may also be in $A^3 \times^4 f^2$ we must have

$$b_0(g + h\alpha - 2l\beta) = 0.$$

If $b_0 = 0$, the intersection is the contact of a tangent plane in the double point; there are six such. If $b_0 \neq 0$, let us take another intersection which is not one of these contacts as $(0, 0, 1)$. Then

$$c = 0,$$

$$b_0(g + h\alpha - 2l\beta) = 0,$$

$$c_0(h + g\alpha - 2i\beta) = 0.$$

These last two equations say that the tangents at the double point, given by

$$x_1^2 + 2lx_1x_2 + fx_2^2 = 0,$$

are of the two sheets respectively in which x_1, c is the polar line

of the algebra with respect to the grading g is

$$x_0 g_2^3 - 3uv g_1 g_2 + 3v g_1^2 g_2 - c g_1^3 = 0.$$

But it is not true that x_0 is the flex line of the plane curve.

This result can be verified by starting with

$$a = a_1 = a_2 = 0$$

and then in g the principal is u, v, w ,

$$3h_0 x_1^2 + 3u_0 x_1 x_2 + 3g_0 x_0 x_2 + b_0 x_1^3 + 3v x_1^2 x_2 + 3u x_1 x_2^2 + c_0 x_2^3,$$

into the equation for w by

$$h - g = w = 0.$$

Then x_0 is the flex line. It meets $A^3 x_f^2$ in g at g

$$-b^2 (5x_1 + 4x_2 x_1^3 x_2 + 3g x_1^2 x_2 + 4c_1 x_1 x_2^3 + 4x_2^4) = 0,$$

while it is the secant line to those in w is x_0 , meets the

quintic. The equation for w in g is the secant line of $A^3 x_f^2$

and x_0 is the flex line of the plane curve g .

This result can be verified algebraically by

$$A^3 x_f^2 = gh \sim 12u_0 x_1^2 + 12c_1 x_1 x_2 + 3g_0 x_0 x_2 + 4u_0 x_0 x_1 + 12v_0 x_0 x_1 x_2$$

$$+ 12v_0 x_0 x_1 x_2 + 4c_0 x_1 x_2 + 4x_1^3 + 3u_1 x_1^2 x_2 + 3g_1 x_1 x_2^2$$

$$+ 4c_1 x_1 x_2^2 + c_0 x_2^3,$$

$$= [3(4x_0x_1^2 + 6x_0x_1x_2 + 3q x_0x_2 + x_1 + 3v x_1^2x_2 - 3v x_1x_2 + c x_2,$$

$$x - 4(gh^2x_0 + 4_0q + 2v - 2c)x_1 + (c_0 + 4q - 2v)x_2]$$

$$= (qv - v^2)x_1^2 - 2v^3x_2) \cdot f(x)$$

It is easy to see that for $x = 0$, the function is zero. For $x \neq 0$, the function is not zero. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$.

The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$.

$$x_1 - x_2 = 0.$$

$$x_1 + x_2 = 1, c, \dots$$

$$x_1 + x_2 = 1, c, \dots$$

$$= x_1 + (v - 1)x_0 + 5q + 2v - 2v)x_1 + c_0 + 4q - 2v)x_2,$$

The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$.

$$A^3 x + f^2 x$$

The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$.

The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$. The function is not zero for $x \neq 0$, and it is not zero for $x = 0$.

3

$$- (x_4)^3 \cdot x_4 \cdot A^{3 \times 2} f^4 + (x_4 \cdot x_4)^3 \cdot A^{3 \times 4} f^2$$

$$- 3 \cdot (x_4)^4 \cdot (x_4)^5 \cdot f^4 = 0$$

Note 1.

$$H = \sum_{i,j,k} \frac{6!}{i!j!k!} H_{ijk}, \quad x_0, x_1, x_2, \dots, x_{j+1}, \dots, x_6,$$

$$H_{600} = agh - al^2 - a_1^2 g - a_2^2 h + 2a, a_2 l,$$

$$H_{510} = \frac{1}{3} (ab_0 g + ahn - 2alm - a_2^2 b_0 - a_1^2 n + 2a, a_2 m \\ - a, gh + a, l^2),$$

$$H_{501} = \frac{1}{3} (ac_0 h + agm - 2aln - a_1^2 c_0 + 2a, a_2 n - a_2^2 m \\ - a_2 gh + a_2 l^2),$$

$$H_{420} = \frac{1}{15} (abg - ba_2^2 + 4ab_0 n - 2ab_2 l + afh - 4am^2 \\ + 2a, a_2 b_2 - a_1^2 f + 2a, b_0 g - 6a_2 b_0 l - 4a, h n \\ + 6a_2 h m + 2a, l m - 3gh^2 + 3hl^2),$$

$$H_{402} = \frac{1}{15} (ach - ca_1^2 + 4ac_0 m - 2ac, l + afg - 4an^2 \\ + 2a, a_2 c, - a_2^2 f + 2a_2 c_0 h - 6a, c_0 l + 6a, g n \\ - 4a_2 g m + 2a_2 l n - 3g^2 h + 3gl^2),$$

$$H_{411} = \frac{1}{15} (2ab_0 c_0 + ab_2 g + ac, h - 2afl - 2am n - a_1^2 c, - a_2^2 b_2 \\ + 2a, a_2 f - 2a, c_0 h - 2a_2 b_0 g + 4a, g n \\ + 4a_2 h n - 2a, l n - 2a_2 l m - 3ghl + 3l^3),$$

$$H_{330} = \frac{1}{10} (abn + ab_0 f + ba, g - 2ab_2 m - 2ba_2 l + 2a_2 b_2 h$$

$$+ 2a_1 b_0 n - a_1 f h - b_0 g h - 2a_1 m^2 - 2b_0 l^2 \\ - 3h^2 n + 6 h l \cdot n),$$

$$H_{321} = \frac{1}{30} (abc_0 + 2ab_0 c_1 - 6a_2 g - 3afn + 2a_1 b_0 c_0 + 4a_1 b_2 g \\ - 2a_1 c_1 h + 2a_2 b_0 n - 4a_2 b_2 l + 5a_2 f h \\ - 3c_0 h^2 - 2a_1 f l - 6b_0 g l - 2a_1 m n - 2a_2 m^2 \\ + 3ghm + 6l^2 n),$$

$$H_{312} = \frac{1}{30} (acb_0 + 2ab_2 c_0 - ca_1 h - 3afn + 2a_2 b_0 c_0 - 2a_2 b_2 g \\ + 4a_2 c_1 h + 2a_1 c_0 n - 4a_1 c_1 l + 5a_1 f g \\ - 3b_0 g^2 - 2a_2 f l - 6c_0 h l - 2a_1 n^2 - 2a_2 m n \\ + 3ghn + 6l^2 n),$$

$$H_{222} = \frac{1}{90} (abc + 2ab_2 c_1 + 2ba_2 c_0 + 2ca_1 b_0 - 3af^2 - 3bg^2 - 3ch^2 \\ + 10a_1 b_2 c_0 + 10a_2 b_0 c_1 - 6a_1 c_1 n - 6a_2 b_2 n \\ - 6b_0 c_0 l - 6a_1 f n - 6a_2 f m - 6b_0 g n \\ - 6b_2 g l - 6c_0 h m - 6c_1 h l + 18fgh + 18lmn).$$

The other coefficients may be obtained from these by

symmetry.

(Vier 2.

$$\tilde{J} = \sum_{i!j!k!} \frac{4!}{i!j!k!} \tilde{J}_{i+j+k} x_0^i x_1^j x_2^k, \quad i+j+k=4,$$

hence

$$\begin{aligned} \tilde{J}_{4,0} = & a v_0 c_0 l - a b_0 g v - a c_0 v^2 + a g^2 \tilde{J} + a v_0 v^2 - a l m v \\ & - a_1 x_2 b_0 c_0 + a_1^2 c_0 v + a_2^2 b_0 v + a_1 b_0 g^2 + a_2 c_0 v^2 \\ & - a_1 c_0 v^2 - a_2 v_0 g l^2 - a_1^2 v^2 + a_1 x_2 v v - a_2^2 v^2 \\ & + a_1 g h v + a_2 g^2 v^2 - 3 a_1 g l v - 3 a_2 b l v \\ & + 2 a_1 l^2 v + 2 a_2 l^2 v - g^2 h^2 + 2 g h l^2 - l^4, \end{aligned}$$

$$\begin{aligned} \tilde{J}_{3,0} = & \frac{1}{4} (2 v c_0 v - a v v^2 - 2 a_1 x_2 c_0 - a_1^2 c_0 v + v_0 v^2 + 2 a_2^2 v \\ & - 2 v_0 f g + b a_1 g^2 + 2 a b_2 g m - a c_1 h m - b a_2 g l \\ & - a b_2 l v + 2 a f h v - a f l m + a_1^2 b_2 c_0 - a_1 a_2 b_0 c_1 \\ & + a_2^2 b_0 f + a_2 b_0 c_0 h + a_1^2 c_1 m + a_1 a_2 b_2 v - 2 a_2^2 b_2 v \\ & - a_1 b_0 c_0 l + a_2 b_2 g h + a_2 c_1 h^2 - 2 a_1^2 f v + a_1 a_2 f m \\ & + 2 a_1 b_0 g v - 3 a_1 b_2 g l - a_1 c_1 h l + 2 a_2 b_2 l^2 - 2 a_2 b_0 l v \\ & + a_1 f g h - b_0 g^2 h - 3 a_2 f h l + 2 a_1 f l^2 - 2 a_1 g v^2 \\ & - a_2 h m v + b_0 g l^2 + a_1 l m v + 2 a_2 l v^2 - g h^2 v \\ & + 2 g h l v + h l^2 v - 2 l^3 v), \end{aligned}$$

$$\tilde{J}_{3,0} = \frac{1}{4} (a c b_0 l - a c h m - c a_1 a_2 b_0 - a b_0 c_1 g + a b_2 c_0 l + c a_1^2 v$$

$$\begin{aligned}
& -ac_0ftv + ca_2bv^2 - afx_2g\ v + 2ac_1\ v - ca_1\ v \\
& - 2c_1b\ v + 2afg\ v - afb\ v - a_1a_2b_2c_0 + a_2''c_0c_1 \\
& + a_1^2c_0f + a_1v_0c_0g - 2a_1^2v_1\ v + 2a_2c_1\ v + 2v_2^2v_3\ v \\
& - a_2v_0c_0b + a_1a_2g^2 + a_1c_1\ v + a_1a_2f\ v - 2a_2^2f\ v \\
& - a_2^2v_2gl + 2a_2c_0v\ v - 3a_2c_1\ b - 2a_1c_0b\ v \\
& + 2a_1c_1v^2 + a_2v_0v\ v - c_0g^2v^2 - 3a_1f\ gl - a_1g^2\ v\ v \\
& + 2a_2f\ b^2 - 2a_2v\ v^2 + c_0b\ b^2 + 2a_1\ v^2 + a_2c_1\ v\ v \\
& - g^2b\ v\ v + 2g^2b\ v + gl^2\ v - 2\beta\ v)_2
\end{aligned}$$

$$\begin{aligned}
v_{22c} = \frac{1}{6} (& 2bc_0\ v + abc_1b - afx_2g - 2v\ v^2 - av_0v_2c_0 - va_1a_2c_1 + av_2^2g \\
& - av_2c_1bv + va_2^2f - va_2c_0v + av_0f\ v - 2v_2bl \\
& + va_1g\ v - va_2g\ v + 2v_2\ v\ v + va_2b\ v + xg^2\ v \\
& + vg^2\ v - af\ v\ v^2 - v\ b^2 + a_1^2v_2c_1 - a_2^2v_0^2 + a_2v_0^2c_0 \\
& + a_1a_2v_2f + a_1v_2c_0\ v + a_2v_0v_2g + a_2v_0c_1\ v \\
& - a_1v_0c_0\ v - a_1v_0c_1v - a_1^2f^2 - v_0^2g^2 - a_1v_2g\ v \\
& - a_2v_0fv + va_2v_2\ v + 2a_1v_0\ v^2 - 3\beta_1v_2\ v \\
& - 3av_2v_0\ v\ v - a_1\ v\ v - 2v_2\ v\ v - v_0\ v^2 \\
& - 2v_2\ v\ v + a_1v\ v\ v + v_0b\ v\ v - a_1\ v^2 + 2a_2^3 \\
& - v_0\ v^2 + 2a_1v^3 + \dots - \dots - \dots \\
& + 4\beta_2\ v\ v - 3\beta^2\ v^2)_2
\end{aligned}$$

Note 3.

The coefficient of x_0^6 in T is obtained from the expression given by Salmon for the invariant T of the cubic as

$$\begin{aligned}
 & a^2 b_0^2 c_0^2 - 3 a^2 b_0 c_0 m n + 4 a^2 b_0 n^3 + 4 a^2 c_0 n^3 - 3 a^3 n^2 n - 3 a a_1 b_0 c_0^2 n \\
 & - 6 a a_2 b_0^2 c_0 g + 6 a a_1 b_0 c_0 g m + 5 a a_2 b_0 c_0 h n + 12 a a_1 b_0 c_0 l n \\
 & + 12 a a_2 b_0 c_0 l m + 4 a b_0^2 g^3 + 4 a c_0^2 h^3 + 12 a b_0 c_0 g h l + 18 a a_1 c_0 h m n \\
 & + 18 a a_2 b_0 g m n - 12 a a_1 b_0 g n^2 - 12 a a_2 c_0 h m^2 - 20 a b_0 c_0 l^3 \\
 & - 24 a a_1 c_0 l m^2 - 24 a a_2 b_0 l n^2 - 12 a b_0 g^2 h n - 12 a c_0 g h^2 m \\
 & - 24 a b_0 g^2 l m - 24 a c_0 h^2 l n + 6 a a_1 g n^2 n - 12 a a_1 n^3 \\
 & - 12 a a_2 g m^3 + 6 a a_2 h n n^2 + 36 a b_0 g l^2 n + 36 a c_0 h l^2 n \\
 & + 12 a a_1 l n n^2 + 12 a a_2 l m^2 n + 24 a g^2 h n^2 + 24 a g h^2 n^2 \\
 & - 60 a g h m n + 12 a g l n^2 + 12 a h l^2 n^2 - 12 a l^3 n n \\
 & + 4 a_1^3 b_0 c_0^2 + 4 a_2^3 b_0^2 c_0 - 12 a_1^2 a_2 b_0 c_0 n - 12 a_1 a_2^2 b_0 c_0 n \\
 & + 18 a_1 a_2 b_0 c_0 g h - 3 a_1^2 c_0^2 n^2 - 3 a_2^2 b_0^2 g^2 - 24 a_1^2 b_0 c_0 g^2 \\
 & - 24 a_2^2 b_0 c_0 l^2 - 12 a_1^2 c_0 n n + 24 a_1^2 a_2 c_0^2 + 24 a_1 a_2 b_0^2 \\
 & - 12 a_2^2 b_0 n n + 36 a_1 a_2 b_0 c_0^2 + 24 a_1^2 b_0^2 c_0^2 + 5 a_1^2 c_0^2 \\
 & + 5 a_1 a_2 b_0^2 + 12 a_1 a_2 c_0^2 + 5 a_2^2 b_0^2 + 2 a_2^2 c_0^2 \\
 & + 12 a_1^2 c_0^2 + 5 a_1 a_2 b_0^2 + 5 a_1 a_2 c_0^2 + 12 a_2^2 b_0^2
 \end{aligned}$$

$$\begin{aligned}
& +12a_1c_0^2v + 12a_2c_0^2v + 6a_1^3 - 12a_1a_2 \\
& -12a_1a_2^2v + a_2^3 - 12a_1c_0c_1^2v - 12a_2c_0c_1^2v + 12a_1c_0^2v \\
& +12a_2c_1^2v - 24a_1^2q^2v - 12a_1^2ql^2v + a_1a_2q^2v \\
& +12a_2q^2v - 24a_2^2q^2v + 12a_1c_0c_1^2v + 12a_2c_0c_1^2v \\
& +36a_1^2ql^2v + 36a_1a_2ql^2v + 36a_1a_2^2ql^2v + 36a_2^2ql^2v \\
& -12a_1c_0^2v - 12a_2c_0^2ql^2 - 24a_1^2q^2v - 12a_1a_2q^2v - 24a_2^2q^2v \\
& -12a_1q^2v - 12a_2q^2v + 36a_1^2ql^2v + 36a_2q^2v \\
& -12a_1qa^2v - 12a_2ql^2v - 36a_1ql^3v - 36a_2ql^3v \\
& +24a_1l^4v + 24a_2l^4v + 8q^3v - 24q^2v^2 + 24ql^4 - 8l^6.
\end{aligned}$$

The other coefficients are too difficult to display in full for the general quartic. They may be obtained in specific cases from the coefficient γ in above or the corresponding coefficient in the differentials.

Note 4

Certain invariants of the quartic have a well-defined geometrical meaning. The two simplest invariants,

the A^3 and A^6 have apolarity meanings and may be included in the list by courtesy. Then comes the A^{15} of Dr. Coble*, which is the condition that the quartic be reducible to the sum of the fourth powers of the six lines of a complete quadrilateral or that the covariant S become 40 conics; the discriminant, A^{27} ; the A^{48} of Dr. Thomassen, expressing that there be a polar conic which is the square of a line; an A^{54} of Dr. Morley's, expressing the condition that the quartic pass through the vertices of a tetrahedron; the

undulation condition, A^{60} . In this list certain gaps are filled up by the A^{45} , the condition that a polar cubic be made up of three lines, and the A^{51} , the condition that the polar cubic be of two points and a square of a line.

There should also be an A^{24} , the condition that the

* Ann. Amer. Math., vol. XXXI, p. 357 (1909)

as the other f as $1 + \dots + t^{20} + \dots$
 For the f $A^{24}f' = 0$
 $A^{24}f'$ for $t = 0$ is 0 $A^{24}f' = 0$
 The f is containing only the first two terms of f^{24}
 $d(t)^{24}$ should be the same as those of f is 0
 multiplied by A^{24} to bring the coefficient of f of f
 degree. If this A^{24} is 0 , then Σ has the same f
 as the other f .

**

Proc. Nat. Ac. Sci., Vol. 3, p. 449 (1917).

Nota

Born February 14, 1892, daughter of Benjamin and
Rebecca Hirschman Cohen. Graduated from Friends
School, Baltimore, Md., in 1909. A.B., Towson College, 1912;
residence at Friends, 1912-13. A.B., Towson College, 1914;
1915; M.A. of Teachers, 1915-16, Friends, 1917-18.

It is with pleasure that I take this opportunity of expressing
my deep gratitude to Professor Morley for his unfailing in-
spiration and guidance throughout my university career. My
thanks and sincere appreciation are also due to Professor
Goble and Professor Green for the instruction and help,
and to Professor Bacon and Professor Lewis of Towson
College for the splendid foundation they laid for my university
studies and for their kindly interest in my work.

Teresa Cohen